




English Teachers On Call

Panic Attacks and Panic Disorder



Copyright © 2018 Surely work.co

完全版テキストはレッスン前に担当講師から受け取って下さい

講師のスカイプチャットにテキスト名を送って下さい

Your teacher can send you the complete material.

Please ask them to send the complete version of this material before the lesson.

attacks may not require treatment. Panic disorder is treated with drug therapy, psychotherapy (eg, exposure therapy, cognitive-behavioral therapy), or both.

Panic attacks are common, affecting as many as 10% of the population in a single year. Most people recover without treatment; a few develop panic disorder. Panic disorder is uncommon, affecting 2 to 3% of the population in a 12-mo period. Panic disorder usually begins in late adolescence or early adulthood and affects women 2 to 3 times more often than men.

Symptoms and Signs

A panic attack involves the sudden onset of at least 4 of the 13 symptoms listed in Table 3: [Anxiety Disorders: Symptoms of a Panic Attack](#). Symptoms usually peak within 10 min and **dissipate** within minutes thereafter, leaving little for a physician to observe. Although uncomfortable—at times extremely so—panic attacks are not medically dangerous.



Copyright © 2018 Surely work.co

完全版テキストはレッスン前に担当講師から受け取って下さい

講師のスカイプチャットにテキスト名を送って下さい

Your teacher can send you the complete material.

Please ask them to send the complete version of this material before the lesson.

panicked. People with panic disorder often worry that they have a dangerous heart, lung, or brain disorder and repeatedly visit their family physician or an emergency department seeking help. Unfortunately, in these settings, attention is often focused on general medical symptoms, and the correct diagnosis sometimes is not made. Many people with panic disorder also have symptoms of major depression.

Table 3

Symptoms of a Panic Attack

Cognitive

Fear of dying

Fear of going crazy or of losing control

Feelings of unreality, strangeness, or detachment from the self (**depersonalization**)

Somatic

Chest pain or discomfort
Dizziness, unsteady feelings, or faintness
Feeling of choking
Flushes or chills
Nausea or abdominal distress
Numbness or tingling sensations
Palpitations or accelerated heart rate
Sensations of shortness of breath or **smothering**
Sweating
Trembling or shaking

Diagnosis

Panic disorder is diagnosed after physical disorders that can mimic anxiety are eliminated and symptoms meet diagnostic criteria **stipulated** in the *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition Text Revision (DSM-IV-TR)*.

Treatment

- Often antidepressants, benzodiazepines, or both
- Often nondrug measures (eg, exposure therapy, cognitive-behavioral therapy)

Some people recover without treatment, particularly if they continue to confront situations in which attacks have occurred. For others, especially without treatment, panic disorder follows a chronic waxing and waning course.

Patients should be told that treatment usually helps control symptoms. If avoidance behaviors have not developed, reassurance, education about anxiety, and encouragement to continue to return to and remain in places where panic attacks have occurred may be all that is needed. However, with a long-standing disorder that involves frequent attacks and avoidance behaviors, treatment is likely to require drug therapy combined with more intensive psychotherapy.

Many drugs can prevent or greatly reduce anticipatory anxiety, phobic avoidance, and the number and intensity of panic attacks:

- **Antidepressants:** The different classes—SSRIs, serotonin-norepinephrine reuptake inhibitors (SNRIs), serotonin modulators, tricyclics (TCAs), and monoamine oxidase inhibitors (MAOIs)—are similarly effective. However, SSRIs and SNRIs offer a potential advantage of fewer adverse effects in comparison with other antidepressants.

- **Benzodiazepines:** These anxiolytics—(see Table 2: [Anxiety Disorders: Benzodiazepines](#)) work more rapidly than antidepressants but are more likely to cause physical dependence and such adverse effects as **somnolence**, **ataxia**, and memory problems. For some patients, long-term use of benzodiazepines is the only effective treatment.
- **Antidepressants plus benzodiazepines:** These drugs are sometimes used in combination initially; the benzodiazepine is slowly tapered after the antidepressant becomes effective (although some patients respond only to the combination treatment).

Panic attacks often recur when drugs are stopped.

Different forms of psychotherapy are effective. Exposure therapy, in which patients confront their fears, helps **diminish** the fear and complications caused by fearful avoidance. For example, patients who fear that they will faint during a panic attack are asked to spin in a chair or to **hyperventilate** until they feel dizzy or faint, thereby learning that they will not faint during an attack. Cognitive-behavioral therapy involves teaching patients to recognize and control their **distorted thinking** and false beliefs and to modify their behavior so that it is more adaptive. For example, if patients describe acceleration of their heart rate or shortness of breath in certain situations or places and fear that they are having a heart attack, they are taught the following:

- Not to avoid those situations
- To understand that their worries are unfounded
- To respond instead with slow, controlled breathing or other methods that promote relaxation

Reference: <http://www.merckmanuals.com>



English Teachers On Call
